


SS3011

SS3011 Space Technology and Applications


“Orbitology” (cont'd)



Naval Postgraduate School
Monterey, California

SS3011


Kepler's First Law: *In a two body universe,
the orbit of a satellite around the Earth is an ellipse with
the Earth centered at one of the focii*




Naval Postgraduate School
Monterey, California

SS3011 Kepler's Second Law

Kepler's Second Law: *In a two body universe, radius vector from the sun (Earth) to the planet (satellite) sweeps out equal areas in equal times*



SS3011 Incremental Area Swept Out by Elliptical Arc



Area Swept out by an Elliptical Arc (cont'd)

$$A_{\text{ellip. arc}} = \int_{v_0}^{v_1} \left[\frac{1}{2} r(v)^2 dv \right] =$$

$$A_{\text{ellip. arc}} = \int_{v_0}^{v_1} \left[\frac{1}{2} \left[\frac{a[1-e^2]}{[1+e \cos(v)]} \right]^2 dv \right] =$$

$$\frac{1}{2} [a[1-e^2]]^2 \int_{v_0}^{v_1} \left[\frac{1}{[1+e \cos(v)]^2} dv \right]$$

"very difficult" integral

Naval Postgraduate School
Monterey, California

Area Swept out by an Elliptical Arc (concluded)

$$A_{\text{ellip. arc}} = \int_{v_0}^{v_1} \left[\frac{1}{2} r(v)^2 dv \right] =$$

Ouch!

$$\frac{1}{2} [a[1-e^2]]^2 \left[\frac{e\sqrt{e^2-1} \sin(v_1) - 2F_1 - 2e \cos(v_1) F_1}{(e^2-1)^{3/2} [1+e \cos(v_1)]} - \frac{e\sqrt{e^2-1} \sin(v_0) - 2F_0 - 2e \cos(v_0) F_0}{(e^2-1)^{3/2} [1+e \cos(v_0)]} \right]$$

⇓

$$F_1 = \tanh^{-1} \left[\frac{(e-1) \tan \left[\frac{v_1}{2} \right]}{\sqrt{e^2-1}} \right] \quad F_0 = \tanh^{-1} \left[\frac{(e-1) \tan \left[\frac{v_0}{2} \right]}{\sqrt{e^2-1}} \right]$$


Naval Postgraduate School
Monterey, California

Total Area of an Elliptical Orbit

Butwe can solve for the total area of an ellipse in closed form

$$A_{\text{ellipse total}} = \int_0^{2\pi} \left[\frac{1}{2} r(v)^2 dv \right] =$$

$$a^2 \pi \sqrt{1 - e^2}$$



$$e = \sqrt{1 - \left[\frac{b}{a}\right]^2} \Rightarrow \sqrt{1 - e^2} = \frac{b}{a} \Rightarrow$$

$$A_{\text{ellipse total}} = a^2 \pi \sqrt{1 - e^2} = \pi a^2 \frac{b}{a} = \boxed{\pi a b}$$

Naval Postgraduate School
Monterey, California

Review: Using Kepler's Second Law to Determine Orbital Position

$$\frac{A_{v_1 - v_0}}{A_{\text{ellipse}}} = \frac{t_1 - t_0}{T} \Rightarrow A_{v_1 - v_0} = \left[a^2 \pi \sqrt{1 - e^2} \right] \frac{t_1 - t_0}{T}$$

$$A_{v_1 - v_0} = A_{v_1 - 0} - A_{v_0 - 0}$$


\Downarrow

$$\boxed{\frac{A_{v_1}}{a^2} = \frac{A_{v_0}}{a^2} + \left[\pi \sqrt{1 - e^2} \right] \frac{t_1 - t_0}{T}}$$

Naval Postgraduate School
Monterey, California

SS3011

Mathematical Representation of Kepler's Second Law



$$\frac{A_{t_2 - t_1}}{A_{\text{ellipse total}}} = \frac{t_2 - t_1}{T} \Rightarrow$$

$$\frac{A_{t_2 - t_1}}{[a^2 \pi \sqrt{1 - e^2}]} = \frac{t_2 - t_1}{T}$$

$$A_{t_2 - t_1} = [a^2 \pi \sqrt{1 - e^2}] \times \frac{t_2 - t_1}{T}$$

School


n i a

SS3011

Review: Graphical Representation of Kepler's Second Law

Graphical Representation of Kepler's Second Law



Or



$$\frac{A_{v_1}}{a^2} = \frac{A_{v_0}}{a^2} + [\pi \sqrt{1 - e^2}] \frac{t_1 - t_0}{T}$$

Graduate School

Monterey, California



Kepler's Third law

- **Kepler's Third Law:** Kepler's Third law *In a two body universe, square of the period of any object revolving about the sun (Earth) is in the same ratio as the cube of its mean distance*

$$\text{constant} = \frac{4a^3 \pi^2}{T^2}$$

$$\mu \equiv \text{constant}$$

Kepler's Third Law

Solving for the
Orbit Period


$$T = \frac{2 \pi a^{3/2}}{\sqrt{\mu}}$$

- Third law directly derivable from first and second laws

$\mu \rightarrow$ **Planetary gravitational parameter**

Naval Postgraduate School
Monterey, California

What is μ ?



Isaac Newton, (1642-1727)

Naval Postgraduate School
Monterey, California

SS3011

Planetary Gravitational Parameter

$$\mu_{\text{earth}} = G M \approx 6.672 \times 10^{-11} \frac{\text{Nt-m}^2}{\text{kg}^2} \times 5.974 \times 10^{24} \text{kg} =$$

$$3.98565 \times 10^{14} \frac{\text{Nt-m}^2}{\text{kg}} = 3.986 \times 10^{14} \frac{\text{m}^3}{\text{sec}^2} = 1.4076 \times 10^{16} \frac{\text{ft}^3}{\text{sec}^2}$$

$$\mu_{\text{moon}} = 4.903 \times 10^3 \frac{\text{m}^3}{\text{sec}^2}$$

$$\mu_{\text{sun}} = 1.327 \times 10^{20} \frac{\text{m}^3}{\text{sec}^2}$$

$$\mu_{\text{Mars}} = 4.269 \times 10^4 \frac{\text{m}^3}{\text{sec}^2}$$

Naval Postgraduate School
Monterey, California

SS3011

Elapsed Time Formulae

(Including Kepler's Third Law)

Elapsed Time Formulae
(Including Kepler's Third Law)

Kepler Second Law

Kepler Third Law

$$T = \frac{2 \pi a^{3/2}}{\sqrt{\mu}}$$

Implicit relationship
between t and v


$$\frac{t_{v_1}}{T} = \frac{A_{v_1}}{A_{\text{ellipse}}} =$$

$$\frac{t_{v_1} \sqrt{\mu}}{2 \pi a^{3/2}} = \frac{[A_{v_1} / a^2]}{\pi \sqrt{[1 - e^2]}} =$$

$$\frac{\frac{1}{2} \int_0^{v_1} \left[\frac{1 - e^2}{1 + e \cos(v)} \right]^2 dv}{\pi \sqrt{[1 - e^2]}}$$


Naval Postgraduate School
Monterey, California

Kelper's Second Law, Normalized Time vs. true anomaly, elliptical orbit




SS3011

Time of Flight Graphs (cont'd)




Propagation of Orbital Position

Kepler's Second Law, Normalized Time vs. true anomaly, elliptical orbit



Homework: Kepler's Second Law



Postscript: Alternate Form of Kepler's Second Law (cont'd)

$$\left\{ \begin{array}{l} \text{Let: } [t_2 \Rightarrow t_1] \rightarrow t_2 - t_1 = dt \\ \text{Then: } A_{t_2 - t_1} = dA(t) \end{array} \right\} \Rightarrow dA(t) = \left[a^2 \pi \sqrt{1-e^2} \right] \frac{dt}{T}$$

• But $dA(t) = \frac{1}{2} r^2 dv$ **"constant" for a given orbit**

and

$$\frac{dA(t)}{dt} = \frac{\left[a^2 \pi \sqrt{1-e^2} \right]}{T} = \frac{\frac{1}{2} r^2 dv}{dt} = \frac{1}{2} r^2 \frac{dv}{dt}$$

"Specific" Angular Momentum

Naval Postgraduate School
Monterey, California


Kepler's Second Law (Concluded)

Alternate form ...

Look at the Problem Physics

- Gravity Can Exert No Torque on a Satellite
- Therefore, Angular momentum is constant

$$r^2 \frac{dv}{dt} = \frac{2 \left[a^2 \pi \sqrt{1-e^2} \right]}{T} \equiv 1$$



Naval Postgraduate School
Monterey, California

Homework: Kepler's Second Law (cont'd)

- United Federation of Planets Starship *Potemtkeon* is parked in a highly elliptical orbit around the Federation Outpost planet *Epsilon-Hydra 5*.
- *Captain Sulu* commanding *Potemtkeon* has orders to *rendezvous* with Federation Starship *Excelsior* commanded by *Captain Checkov*
- From sensor readings *Captain Sulu* knows that the orbit around *Epsilon-Hydra 5* has a closest approach (*pera-Epsilon*) of *0.25 astronomical units* (AU)*. The farthest distance from the planet (*ap-Epsilon*) is *0.75 AU*
- *The Potemtkeon* has passed the orbit *ap-Epsilon* and is headed back towards the closest approach to the planet
- Navigation data shows that the current position of the *Potemtkeon* relative to the planet is EXACTLY PERPENDICULAR to the *Line of Apsides** of the orbit
*(the line connecting the *pera-Epsilon* and the *ap-Epsilon*)

*AU \approx 150,000,000 kilometers

N a v a l P o s t g r a d u a t e S c h o o l
M o n t e r e y , C a l i f o r n i a

Homework: Kepler's Second Law (cont'd)

- The onboard atomic clock shows that it has been exactly *50 hours, 21 minutes, 48 seconds, and 600 milliseconds* (*3021.81 minutes*) since the *Potemtkeon* last passed the *pera-Epsilon* point
- In order to successfully *rendezvous*, *Captain Checkov* aboard the *Excelsior* must know the exact orbital eccentricity, *e*, and the time of arrival at the *pera-Epsilon* point from the current position
- What should *Captain Sulu* tell him?
- If they fail to *rendezvous* at the next encounter, how long will *Checkov* have to wait until the next *pera-Epsilon* encounter

N a v a l P o s t g r a d u a t e S c h o o l
M o n t e r e y , C a l i f o r n i a

Homework: Kepler's Second Law (concluded)

• Hint 1:

$$\begin{bmatrix} r_{\min} = a(1 - e) \\ r_{\max} = a(1 + e) \end{bmatrix}$$

• Hint 2:

• Hint 3:

$$\frac{A\left(\frac{3\pi}{2}\right)}{A_{\text{total}}} = \frac{t\left(\frac{3\pi}{2}\right)}{T}$$

N a v

$$\int_{-\frac{\pi}{2}}^0 \left[\frac{dv}{[1 + e \cos(v)]^2} \right] =$$

$$\left[\frac{1}{1 - e^2} \right] \left[\frac{2 \tan^{-1} \left[\frac{1 - e}{\sqrt{1 - e^2}} \right]}{\sqrt{1 - e^2}} - e \right]$$

Monterey, California

Homework: Kepler's Second Law (concluded)

Hint 4

Homework ...

..... Or Instead of evaluating the BRUTAL Integral

$$\int_{-\frac{\pi}{2}}^0 \left[\frac{dv}{[1 + e \cos(v)]^2} \right] =$$

$$\left[\frac{1}{1 - e^2} \right] \left[\frac{2 \tan^{-1} \left[\frac{1 - e}{\sqrt{1 - e^2}} \right]}{\sqrt{1 - e^2}} - e \right]$$

You can use the area Swept Area or time of flight
graphs for Kepler's second law